

UC Davis

UC Davis Previously Published Works

Title

Tuning the kinetics of cadherin adhesion.

Permalink

<https://escholarship.org/uc/item/61r0418j>

Journal

The Journal of investigative dermatology, 133(10)

ISSN

0022-202X

Author

Sivasankar, Sanjeevi

Publication Date

2013-10-01

DOI

10.1038/jid.2013.229

Peer reviewed



Published in final edited form as:

J Invest Dermatol. 2013 October ; 133(10): 2318–2323. doi:10.1038/jid.2013.229.

Tuning the kinetics of cadherin adhesion

Sanjeevi Sivasankar

Department of Physics and Astronomy, Iowa State University, Ames, IA 50011

Ames Laboratory, United States Department of Energy, Ames, IA 50011

Abstract

Cadherins are Ca^{2+} dependent cell-cell adhesion proteins that maintain the structural integrity of the epidermis; their principle function is to resist mechanical force. This review summarizes the biophysical mechanisms by which classical cadherins tune adhesion and withstand mechanical stress. We first relate the structure of classical cadherins to their equilibrium binding properties. We then review the role of mechanical perturbations in tuning the kinetics of cadherin adhesion. In particular, we highlight recent studies which show that cadherins form three types of adhesive bonds: *catch bonds* which become longer lived and lock in the presence of tensile force, *slip bonds* which become shorter lived when pulled and *ideal bonds* that are insensitive to tugging.

Introduction

The epidermis serves as a physical barrier that protects organisms from their external environment. This multilayered tissue is composed of keratinocytes bound together by two types of cell-cell adhesion complexes: desmosomes and adherens junctions. (Jensen and Wheelock, 1996). The primary adhesive components of both these structures are the cadherin family of Ca^{2+} dependent transmembrane proteins (Al-Amoudi and Frangakis, 2008; Green and Simpson, 2007; Gumbiner, 2005; Halbleib and Nelson, 2006; Niessen *et al.*, 2011). Desmosomes are composed of two types of desmosomal cadherins (desmocollin and desmoglein) (Desai *et al.*, 2009; Green and Simpson, 2007), while epidermal adherens junctions contain a single classical type-1 cadherin (either E-cadherin or P-cadherin) (Halbleib and Nelson, 2006; Jensen and Wheelock, 1996). Both desmosomes and adherens junctions act in a coordinated fashion to help the epidermis withstand mechanical stress. While the interactions that mediate desmosomal cadherin binding are not completely understood, the structural basis of classical cadherin adhesion has been extensively characterized.

Classical cadherins share a conserved cytoplasmic domain, and an ectodomain containing five tandem extracellular (EC) repeats. Their expression levels vary within the epidermis; while E-cadherins are present in all keratinocytes, expression of P-cadherins is limited to the basal layer (Halbleib and Nelson, 2006; Takeichi, 1988). Adhesion is mediated by the

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Corresponding author: Sanjeevi Sivasankar, Tel: (515) 294-1220; Fax: (515) 294-6027; sivasank@iastate.edu.

The author declares no conflict of interest.

cadherin ectodomain while the cytoplasmic region binds to adaptor proteins which link cadherins indirectly to the cytoskeleton, regulate cadherin turnover and modulate actin assembly (Nelson and Nusse, 2004; Niessen *et al.*, 2011; Takeichi, 2007). Since the epidermis is a self-renewing tissue with a continuous upward movement of cells, cadherins dynamically tune their adhesive strength in order to preserve epidermal barrier integrity (Niessen, 2007). Epidermal cadherin knockout studies in mice show that loss of E-cadherin correlates with a loss of adherens junctions, altered epidermal differentiation and loss of hair follicles (Tinkle *et al.*, 2004; Young *et al.*, 2003). Similarly, deletion of α -catenin, an adaptor protein associated with the cadherin cytoplasmic domain, results in impaired adhesion and epidermal detachment (Vasioukhin *et al.*, 2001).

Cell-cell adhesion is a dynamic process and classical cadherins tailor their binding kinetics in order to withstand mechanical perturbations. While the equilibrium binding properties of classical cadherins have been extensively characterized (Brasch *et al.*, 2012), the role of mechanical force in altering cadherin binding is only now being measured. Recent studies show that upon being exposed to mechanical perturbation, E-cadherins change their unbinding kinetics (Rakshit *et al.*, 2012). These kinetic changes are not manifested in solution or in the absence of mechanical loading, but are critical for cadherin adhesion.

This brief review summarizes our current understanding of the effect of mechanical force on the kinetics of E-cadherin adhesion. We focus on the ectodomain; the role of the cytoplasmic domain and its associated proteins have been reviewed elsewhere (Gomez *et al.*, 2011; Ladoux and Nicolas, 2012; Leckband *et al.*, 2011; Papusheva and Heisenberg, 2010; Schwartz and DeSimone, 2008). We begin by relating the structure of E-cadherins to their equilibrium binding properties. We then review the role of mechanical perturbations in tuning the kinetics of adhesion. Finally, we discuss major open questions and future directions in this exciting area of research.

Adhesive states of classical cadherins

Classical cadherins adhere via ‘*trans*’ interactions where ectodomains from opposing cells bridge the inter-membrane gap and interact with each other. Adhesion is strengthened by the cooperative self-assembly of cadherins on the same cell into *cis* clusters (Brasch *et al.*, 2012).

Structure and kinetics of *trans* adhesive states

Structural studies of both the complete ectodomain of type I classical C-cadherin (EC1–5) (Boggon *et al.*, 2002) and of smaller fragments of E-Cadherin and N-Cadherin (Harrison *et al.*, 2010; Haussinger *et al.*, 2004; Nagar *et al.*, 1996; Pertz *et al.*, 1999; Shapiro *et al.*, 1995) have identified key interactions that mediate *trans* adhesion (Fig. 1). The primary adhesive conformation involves the interaction of opposing EC1 domains and is termed the strand-swapped dimer (Fig. 1A). In this structure, N-terminal β -strands between opposing EC1 domains are swapped and the side chain of a conserved Tryptophan at position 2 (W2) is inserted into a pocket on their adhesive partner (Boggon *et al.*, 2002; Haussinger *et al.*, 2004; Parisini *et al.*, 2007; Shapiro *et al.*, 1995) (Fig. 1A). The physiological relevance of this adhesive interface has been confirmed in numerous mutational, structural and cellular

studies (Harrison *et al.*, 2005; Pertz *et al.*, 1999; Prakasam *et al.*, 2006a; Shan *et al.*, 2004; Tamura *et al.*, 1998; Troyanovsky *et al.*, 2003). In solution, the affinity for strand-swap dimer formation is low; dissociation constants (K_d) for *trans* dimers of the full length ectodomain of C-cadherin measured using Analytical Ultra Centrifugation (AUC) is 64 μ M (Chappuis-Flament *et al.*, 2001). Similarly, EC1–2 domains of E-cadherin expressed in mammalian and bacterial cells have *trans* dimer K_d values of 97 μ M (Katsamba *et al.*, 2009) and 80 μ M (Koch *et al.*, 1997) respectively.

Prior to strand-swapping, cadherin monomers are in a “closed” conformation where W2 is docked into each monomer’s binding pocket; the monomers thus act as competitive inhibitors of strand-swapping (Chen *et al.*, 2005). The closed monomeric conformation places a strain on the short swapping strand due to its anchorage at one end by the W2 and at the other by a Ca^{2+} ion; relieving this conformational strain is the driving force for strand swapping (Vendome *et al.*, 2011). Equilibrium affinity measurements using AUC show that mutations that relieve strain in the swapping strand in E-cadherin monomers, decrease dimerization affinities (Vendome *et al.*, 2011). Single molecule Fluorescence Resonance Energy Transfer (FRET) experiments suggest that prior to swapping N-terminal β -strands, E-cadherin monomers first form a non-swapped, intermediate “encounter complex” (Fig. 1C) (Sivasankar *et al.*, 2009). E-cadherins can be trapped in this encounter complex by mutating W2 (Sivasankar *et al.*, 2009); consequently, W2A fragments weakly adhere to each other (Prakasam *et al.*, 2006a; Sivasankar *et al.*, 2009). Recently, the atomic resolution structure of the encounter complex has been resolved in W2A mutants (Fig. 1B). This conformation, called an X-dimer, is formed by extensive surface interactions between the base of the EC1 domain, EC1–EC2 interdomain linker region and the apex of domain EC2 (Harrison *et al.*, 2010) (Fig. 1B). The affinity for X-dimer formation in solution is significantly weaker than strand-swap dimers; the K_d of W2A cadherin X-dimers is an order of magnitude higher (916 μ M) than wild type (WT) cadherin strand-swap dimers (Harrison *et al.*, 2010).

Mutations in the cadherin X-dimer binding interface alter the kinetics of strand-swapping but do not change the structure of the strand-swap dimer. When a key Lys 14 residue in the X-dimer binding interface is mutated to a Glu, the *trans*-dimers are virtually indistinguishable from WT cadherin strand-swap dimers (Harrison *et al.*, 2010). As measured using Surface Plasmon Resonance, the K14E mutants show no binding in a short-time frame suggesting that their binding rate (on rate) is low. Similarly, sedimentation velocity AUC and size-exclusion chromatography show that the monomer to strand-swap dimer conversion is impeded in these mutants (Harrison *et al.*, 2010). Presumably, lower on-rates are measured since the formation of X-dimers, which serve as kinetic intermediates strand-swapping (Fig. 1C), are impaired in the K14E mutants. In epithelial cells, inactivation of X dimers result in extraordinarily stable cell-cell junctions; this has been interpreted to indicate that X-dimers are an intermediate in the pathway to dissociation of strand-swap dimers (Hong *et al.*, 2011). AUC measurements show that the K_d of the K14E mutants are virtually indistinguishable from WT cadherin which suggests that besides their low on-rate, the dissociation (off-rate) of these mutants is also decreased (Harrison *et al.*, 2010). However, in contrast to these studies, recent single molecule force measurements indicate

that the dissociation rate of K14E is similar to WT cadherin (Rakshit *et al.*, 2012). Consequently, the molecular role of X-dimers in the dissociation of strand-swap dimers is unclear.

Structure and kinetics of *cis* adhesive states

Cadherin adhesion is enhanced by their lateral assembly on the cell surface (Kim *et al.*, 2005; Takeda *et al.*, 1999). However, the biophysical mechanisms by which *cis* clustering boosts adhesion are just beginning to be understood. Early studies showed that beads decorated with cadherin pairs aggregated to a greater extent than beads with immobilized monomers (Briher *et al.*, 1996). While this data was interpreted to suggest that cadherin ectodomains form *cis* dimers, recent single molecule experiments show that ectodomains located adjacent to each cooperatively enhance the probability of adhesion even if they do not associate with each other in a *cis* geometry (Zhang *et al.*, 2009).

Based on contacts observed in X-ray crystal structures of a range of classical cadherins, it has been proposed that interactions between the apex of EC1 and the base of EC2 of neighboring cadherins mediate dimerization in a *cis* orientation (Boggon *et al.*, 2002; Harrison *et al.*, 2011). These interactions are however not observed in NMR measurements of EC1–2 (Haussinger *et al.*, 2002), indicating that their K_d exceeds 1mM (Harrison *et al.*, 2011). Similarly, single molecule FRET experiments could not detect *cis* dimer formation between two cadherin ectodomains that were located adjacent to each other in a configuration that would permit lateral dimerization (Zhang *et al.*, 2009). This discrepancy is explained by recent theoretical studies which predict that the *cis* assembly of cadherin ectodomains requires prior *trans* dimerization (Wu *et al.*, 2011; Wu *et al.*, 2010). When *trans* dimers are formed, the conformational flexibility of ectodomains is dramatically reduced which lowers the entropic penalty associated with *cis* dimer formation (Wu *et al.*, 2011).

In qualitative agreement with these predictions, micropipette manipulation experiments show that the binding of cadherins from opposing cells occur in two stages: an initial rapid stage ascribed to *trans* adhesion followed by a second, slower stage interpreted to occur due to *cis* clustering (Chien *et al.*, 2008). However, while the first stage requires the EC1 domain as expected for *trans* dimer formation, EC3 is required for the second adhesive state (Chien *et al.*, 2008). Micropipette experiments also demonstrate that hypoglycosylation of EC2 and EC3 enhance the lateral assembly of ectodomains (Langer *et al.*, 2012).

Mechanical tension alters the kinetics of cadherin adhesion

The structural and biophysical studies described above, provide a detailed picture of the kinetic determinants of classical cadherin binding in equilibrium, under force-free conditions. However, the molecular mechanisms by which cadherins alter their binding kinetics in response to mechanical forces are still unclear.

When cadherin *trans* dimers, are pulled apart, they can form one of three distinct types of bonds (Dembo, 1994; Dembo *et al.*, 1988) (i) *Slip bonds* which weaken and have a higher off-rate when pulled. (ii) *Catch bonds* which counter-intuitively strengthen such that their

off-rates decrease. (iii) *Ideal bonds* which are unaffected by mechanical stress. Slip bonds are the most commonly observed interactions in biology. Catch bonds provide a way for the interacting proteins to grip tightly in the presence of tugging forces. Finally, though ideal bonds were theoretically proposed more than a decade ago (Dembo, 1994; Dembo *et al.*, 1988), they had not been experimentally observed in any biological system.

Recently, single molecule Atomic Force Microscope (AFM) force measurements were used to show that E-cadherins form bonds with catch, slip and ideal mechanical properties (Rakshit *et al.*, 2012). The lifetimes of E-cadherin binding conformations were measured as they were subjected to different pulling forces. These experiments showed that while W2A mutant X-dimers formed catch bonds, WT and K14E strand-swap dimers formed slip bonds (Rakshit *et al.*, 2012) (Fig. 2A & 2B). WT cadherins were also shown to form ideal bonds which were hypothesized to arise as X-dimers converted to a strand-swap conformation (Fig. 2B) (Rakshit *et al.*, 2012).

X-dimers form catch bonds

When X-dimers were tugged, their bond lifetimes increased with force, indicative of a catch bond. After reaching a maximum at a critical force of ~ 30 pN, the lifetimes decreased with force (Fig. 2A). A similar behavior was measured when WT cadherins were forced into an X-dimer conformation by competitively inhibiting strand swapping using free W in solution (Rakshit *et al.*, 2012). X-dimer catch bonds are observed because the cadherins reorient when they are pulled such that they form transient, force-induced bonds and lock more tightly.

While this was the first observation of catch bonds in cadherin adhesion, these bonds have previously been measured with other adhesive proteins like selectins (Marshall *et al.*, 2003; Sarangapani *et al.*, 2004; Yago *et al.*, 2004), FimH (Le Trong *et al.*, 2010; Thomas *et al.*, 2002) and integrins (Kong *et al.*, 2009). Although it is tempting to speculate that the physiological role of X-dimer catch bonds is to allow cells to grip tightly and lock in place when pulled; this hypothesis remains to be tested.

Catch bonds resolve discrepancies between solution and surface force measurements

Over a decade ago, Surface Force Apparatus (SFA) measurements of the interactions between cadherin ectodomains immobilized on lipid membranes suggested that classical cadherins bind in three distinct conformations (Sivasankar *et al.*, 2001). The weakest conformation required W2, and corresponded to a strand swapped dimer (Prakasam *et al.*, 2006b; Zhu *et al.*, 2003). The second conformation had an intermediate binding strength and required EC1–2 (Zhu *et al.*, 2003); based on recent structural data, it is likely that this adhesive state corresponds to the X-dimer complex. The third and strongest adhesion required the EC3 domains to interact directly (Zhu *et al.*, 2003); while this adhesive state likely corresponds to a *cis*-dimer structure, this remains to be confirmed. Single molecule AFM force measurements of the interaction of different classical cadherins confirmed the results of the ensemble SFA measurements (Bayas *et al.*, 2006; Perret *et al.*, 2004; Shi *et al.*, 2008; Shi *et al.*, 2010).

It was initially believed that the SFA measurement of stronger adhesion between X-dimers compared to strand-swap dimers directly contradicted the results of solution affinity measurements which showed that X-dimers have higher off rates than strand-swap dimers. However, the discovery of X-dimer catch bonds resolves this apparent discrepancy (Fig. 2A). Since catch bonds strengthen in the presence of force, X-dimer adhesion which is weak in the absence of force becomes stronger when pulled.

Strand swap dimers form slip bonds

Since strand-swap dimers have a higher binding affinity than X-dimers (Harrison *et al.*, 2010; Katsamba *et al.*, 2009), WT cadherins form strand-swap dimers when they interact for long periods of time. Single molecule AFM force clamp experiments showed that these WT cadherin strand-swap dimers formed slip bonds; their bond lifetimes decreased with increasing tensile force (Fig. 2B). Not surprisingly, identical slip bonds were formed by the K14E strand-swap dimers (Rakshit *et al.*, 2012) (Fig. 2B). The intrinsic off rate of both the WT E-cadherin and K14E strand swap dimers was 1.6 s^{-1} (Rakshit *et al.*, 2012) which is similar to an off-rate of 0.7 s^{-1} measured for WT E-cadherins using NMR (Haussinger *et al.*, 2004).

Ideal bonds are formed as X-dimers transition to a strand-swap conformation

Besides forming catch and slip bonds, cadherins also form ideal bonds that behave like mechanical dampers and prevent the abrupt jolting of cells. When WT cadherin interaction time was decreased, the lifetimes of their interactions were independent of force; they formed ideal bonds (Rakshit *et al.*, 2012) (Fig. 2B). It was hypothesized that ideal bonds correspond to an intermediate state which is formed when X-dimers transition to strand-swap binding (Rakshit *et al.*, 2012). However, the structure of the intermediate state and the molecular contacts responsible for ideal bond formation still need to be resolved.

Future Directions

Catch, slip and ideal bonds suggest a physical mechanism that E-cadherins use to resist tensile force as cells rearrange during skin renewal and wound healing. It is tempting to speculate that as keratinocytes reposition themselves, E-cadherins bind rapidly to form X-dimer catch bonds that allow cells to grip strongly under load (Rakshit *et al.*, 2012) (Fig. 2C). Over time, the X-dimers proceed to form more robust strand-swap dimers that have a high affinity in the absence of force (Fig. 2C); this conversion is facilitated by an intermediate conformation that is insensitive to tensile force (Rakshit *et al.*, 2012). However, it is currently unclear if keratinocytes utilize such a mechanism to tune adhesive properties. Studying cadherin bond mechanics in living cells will be a crucial first step to addressing this question.

Besides mediating robust adhesion, classical cadherins play a key role in mechanotransduction by sensing physical stimuli at cell-cell junctions, transmitting them to the cytoplasm and activating a biochemical response (Ladoux *et al.*, 2010; le Duc *et al.*, 2010; Liu *et al.*, 2010; Weber *et al.*, 2012). It is believed that cadherins along with their adaptor proteins, β -catenin and α -catenin form the core force-bearing unit in the

transmission of mechanical signals (Leckband *et al.*, 2011). To accomplish this, it is critical that the interactions between cadherins and catenins remain intact when exposed to force. However, the force dependent binding kinetics of these interactions have not yet been studied. Furthermore, the role of these adapter proteins in altering cadherin mechanical properties is still an open question. For instance, it is known that the adapter protein α -catenin plays an important role in strengthening cadherin bonds following initial adhesion (Bajpai *et al.*, 2008). Whether α -catenin and other adapter proteins alter the force dependent kinetics of cadherin bonds needs to be investigated.

Some of the discrepancies in cadherin binding measured using solution affinity measurements and force measurements arise due to differences between cadherin interactions in solution, under force independent conditions, and cadherin adhesion in the presence of mechanical stress. The discovery that cadherins vary their lifetimes in response to force reconciles some of these differences (Rakshit *et al.*, 2012). However, several open questions remain. For instance, the molecular interactions by which cadherins form catch bonds are not known. Furthermore, the hypothesis that ideal bonds correspond to an intermediate state which is formed as cadherin X-dimers transition to a strand-swap conformation needs to be tested at the molecular level. Finally, the role that X-dimers play in the dissociation of strand-swap dimers is unclear.

Acknowledgments

The author thanks Dr. Sabyasachi Rakshit for his help in preparing Figure 2. This work was supported in part by a Basil O'Connor Starter Scholar Award from the March of Dimes Foundation and an American Heart Association National Scientist Development Grant.

References

- Al-Amoudi A, Frangakis AS. Structural studies on desmosomes. *Biochem Soc Trans.* 2008; 36:181–187. [PubMed: 18363559]
- Bajpai S, Correia J, Feng YF, Figueiredo J, Sun SX, Longmore GD, et al. α -Catenin mediates initial E-cadherin-dependent cell-cell recognition and subsequent bond strengthening. *Proc Natl Acad Sci U S A.* 2008; 105:18331–18336. [PubMed: 19017792]
- Bayas MV, Leung A, Evans E, Leckband D. Lifetime measurements reveal kinetic differences between homophilic cadherin bonds. *Biophys J.* 2006; 90:1385–1395. [PubMed: 16326909]
- Boggon TJ, Murray J, Chappuis-Flament S, Wong E, Gumbiner BM, Shapiro L. C-cadherin ectodomain structure and implications for cell adhesion mechanisms. *Science.* 2002; 296:1308–1313. [PubMed: 11964443]
- Brasch J, Harrison OJ, Honig B, Shapiro L. Thinking outside the cell: how cadherins drive adhesion. *Trends Cell Biol.* 2012; 22:299–310. [PubMed: 22555008]
- Brieher WM, Yap AS, Gumbiner BM. Lateral dimerization is required for the homophilic binding activity of C-cadherin. *J Cell Biol.* 1996; 135:487–496. [PubMed: 8896604]
- Chappuis-Flament S, Wong E, Hicks LD, Kay CM, Gumbiner BM. Multiple cadherin extracellular repeats mediate homophilic binding and adhesion. *Journal of Cell Biology.* 2001; 154:231–243. [PubMed: 11449003]
- Chen CP, Posy S, Ben-Shaul A, Shapiro L, Honig BH. Specificity of cell-cell adhesion by classical cadherins: Critical role for low-affinity dimerization through beta-strand swapping. *Proc Natl Acad Sci U S A.* 2005; 102:8531–8536. [PubMed: 15937105]
- Chien YH, Jiang N, Li F, Zhang F, Zhu C, Leckband D. Two stage cadherin kinetics require multiple extracellular domains but not the cytoplasmic region. *J Biol Chem.* 2008; 283:1848–1856. [PubMed: 17999960]

- Dembo, M. Lectures on Mathematics in the Life Sciences, Some Mathematical Problems in Biology. Vol. Vol. 24. Providence, Rhode Island: American Mathematical Society; 1994. On peeling an adherent cell from a surface; p. 51-77.
- Dembo M, Torney DC, Saxman K, Hammer D. The Reaction-Limited Kinetics of Membrane-to-Surface Adhesion and Detachment. *Proc R Soc Lond B Biol Sci.* 1988; 234:55–83. [PubMed: 2901109]
- Desai BV, Harmon RM, Green KJ. Desmosomes at a glance. *J Cell Sci.* 2009; 122:4401–4407. [PubMed: 19955337]
- Gomez GA, McLachlan RW, Yap AS. Productive tension: force-sensing and homeostasis of cell-cell junctions. *Trends Cell Biol.* 2011; 21:499–505. [PubMed: 21763139]
- Green KJ, Simpson CL. Desmosomes: New perspectives on a classic. *J Invest Dermatol.* 2007; 127:2499–2515. [PubMed: 17934502]
- Gumbiner BM. Regulation of cadherin-mediated adhesion in morphogenesis. *Nat Rev Mol Cell Biol.* 2005; 6:622–634. [PubMed: 16025097]
- Halbleib JM, Nelson WJ. Cadherins in development: cell adhesion, sorting, and tissue morphogenesis. *Genes Dev.* 2006; 20:3199–3214. [PubMed: 17158740]
- Harrison OJ, Bahna F, Katsamba PS, Jin XS, Brasch J, Vendome J, et al. Two-step adhesive binding by classical cadherins. *Nat Struct Mol Biol.* 2010; 17 348-U121.
- Harrison OJ, Corps EM, Berge T, Kilshaw PJ. The mechanism of cell adhesion by classical cadherins: the role of domain 1. *J Cell Sci.* 2005; 118:711–721. [PubMed: 15671061]
- Harrison OJ, Jin XS, Hong SJ, Bahna F, Ahlsen G, Brasch J, et al. The Extracellular Architecture of Adherens Junctions Revealed by Crystal Structures of Type I Cadherins. *Structure.* 2011; 19:244–256. [PubMed: 21300292]
- Haussinger D, Ahrens T, Aberle T, Engel J, Stetefeld J, Grzesiek S. Proteolytic E-cadherin activation followed by solution NMR and X-ray crystallography. *EMBO J.* 2004; 23:1699–1708. [PubMed: 15071499]
- Haussinger D, Ahrens T, Sass HJ, Pertz O, Engel J, Grzesiek S. Calcium-dependent homoassociation of E-cadherin by NMR spectroscopy: changes in mobility, conformation and mapping of contact regions. *J Mol Biol.* 2002; 324:823–839. [PubMed: 12460580]
- Hong SJ, Troyanovsky RB, Troyanovsky SM. Cadherin exits the junction by switching its adhesive bond. *J Cell Biol.* 2011; 192:1073–1083. [PubMed: 21422232]
- Jensen PJ, Wheelock MJ. The relationships among adhesion, stratification and differentiation in keratinocytes. *Cell Death and Differentiation.* 1996; 3:357–371. [PubMed: 17180106]
- Katsamba P, Carroll K, Ahlsen G, Bahna F, Vendome J, Posy S, et al. Linking molecular affinity and cellular specificity in cadherin-mediated adhesion. *Proc Natl Acad Sci U S A.* 2009; 106:11594–11599. [PubMed: 19553217]
- Kim YJ, Johnson KR, Wheelock MJ. N-cadherin-mediated cell motility requires cis dimers. *Cell Commun Adhes.* 2005; 12:23–39. [PubMed: 16371344]
- Koch AW, Pokutta S, Lustig A, Engel J. Calcium binding and homoassociation of E-cadherin domains. *Biochemistry.* 1997; 36:7697–7705. [PubMed: 9201910]
- Kong F, Garcia AJ, Mould AP, Humphries MJ, Zhu C. Demonstration of catch bonds between an integrin and its ligand. *J Cell Biol.* 2009; 185:1275–1284. [PubMed: 19564406]
- Ladoux B, Anon E, Lambert M, Rabodzey A, Hersen P, Buguin A, et al. Strength Dependence of Cadherin-Mediated Adhesions. *Biophys J.* 2010; 98:534–542. [PubMed: 20159149]
- Ladoux B, Nicolas A. Physically based principles of cell adhesion mechanosensitivity in tissues. *Reports on Progress in Physics.* 2012; 75
- Langer MD, Guo HB, Shashikanth N, Pierce JM, Leckband DE. N-glycosylation alters cadherin-mediated intercellular binding kinetics. *J Cell Sci.* 2012; 125:2478–2485. [PubMed: 22344255]
- le Duc Q, Shi Q, Blonk I, Sonnenberg A, Wang N, Leckband D, et al. Vinculin potentiates E-cadherin mechanosensing and is recruited to actin-anchored sites within adherens junctions in a myosin II-dependent manner. *J Cell Biol.* 2010; 189:1107–1115. [PubMed: 20584916]

- Le Trong I, Aprikian P, Kidd BA, Forero-Shelton M, Tchesnokova V, Rajagopal P, et al. Structural Basis for Mechanical Force Regulation of the Adhesin FimH via Finger Trap-like beta Sheet Twisting. *Cell*. 2010; 141:645–655. [PubMed: 20478255]
- Leckband DE, le Duc Q, Wang N, de Rooij J. Mechanotransduction at cadherin-mediated adhesions. *Curr Opin Cell Biol*. 2011; 23:523–530. [PubMed: 21890337]
- Liu ZJ, Tan JL, Cohen DM, Yang MT, Sniadecki NJ, Ruiz SA, et al. Mechanical tugging force regulates the size of cell-cell junctions. *Proc Natl Acad Sci U S A*. 2010; 107:9944–9949. [PubMed: 20463286]
- Marshall BT, Long M, Piper JW, Yago T, McEver RP, Zhu C. Direct observation of catch bonds involving cell-adhesion molecules. *Nature*. 2003; 423:190–193. [PubMed: 12736689]
- Nagar B, Overduin M, Ikura M, Rini JM. Structural basis of calcium-induced E-cadherin rigidification and dimerization. *Nature*. 1996; 380:360–364. [PubMed: 8598933]
- Nelson WJ, Nusse R. Convergence of Wnt, beta-catenin, and cadherin pathways. *Science*. 2004; 303:1483–1487. [PubMed: 15001769]
- Niessen CM. Tight junctions/adherens junctions: Basic structure and function. *J Invest Dermatol*. 2007; 127:2525–2532. [PubMed: 17934504]
- Niessen CM, Leckband D, Yap AS. Tissue Organization by Cadherin Adhesion Molecules: Dynamic Molecular and Cellular Mechanisms of Morphogenetic Regulation. *Physiol Rev*. 2011; 91:691–731. [PubMed: 21527735]
- Papusheva E, Heisenberg C-P. Spatial organization of adhesion: force-dependent regulation and function in tissue morphogenesis. *EMBO J*. 2010; 29:2753–2768. [PubMed: 20717145]
- Parisini E, Higgins JMG, Liu JH, Brenner MB, Wang JH. The crystal structure of human E-cadherin domains 1 and 2, and comparison with other cadherins in the context of adhesion mechanism. *J Mol Biol*. 2007; 373:401–411. [PubMed: 17850815]
- Perret E, Leung A, Feracci H, Evans E. Trans-bonded pairs of E-cadherin exhibit a remarkable hierarchy of mechanical strengths. *Proc Natl Acad Sci U S A*. 2004; 101:16472–16477. [PubMed: 15546992]
- Pertz O, Bozic D, Koch AW, Fauser C, Brancaccio A, Engel J. A new crystal structure, Ca²⁺ dependence and mutational analysis reveal molecular details of E-cadherin homoassociation. *EMBO J*. 1999; 18:1738–1747. [PubMed: 10202138]
- Prakasam A, Chien YH, Maruthamuthu V, Leckband DE. Calcium site mutations in cadherin: impact on adhesion and evidence of cooperativity. *Biochemistry*. 2006a; 45:6930–6939. [PubMed: 16734428]
- Prakasam AK, Maruthamuthu V, Leckband DE. Similarities between heterophilic and homophilic cadherin adhesion. *Proc Natl Acad Sci U S A*. 2006b; 103:15434–15439. [PubMed: 17023539]
- Rakshit S, Zhang Y, Manibog K, Shafraz O, Sivasankar S. Ideal, catch, and slip bonds in cadherin adhesion. *Proc Natl Acad Sci U S A*. 2012; 109:18815–18820. [PubMed: 23112161]
- Sarangapani KK, Yago T, Klopocki AG, Lawrence MB, Fieger CB, Rosen SD, et al. Low force decelerates L-selectin dissociation from P-selectin glycoprotein ligand-1 and endoglycan. *J Biol Chem*. 2004; 279:2291–2298. [PubMed: 14573602]
- Schwartz MA, DeSimone DW. Cell adhesion receptors in mechanotransduction. *Curr Opin Cell Biol*. 2008; 20:551–556. [PubMed: 18583124]
- Shan W, Yagita Y, Wang Z, Koch A, Fex Svenningsen A, Gruzglin E, et al. The minimal essential unit for cadherin-mediated intercellular adhesion comprises extracellular domains 1 and 2. *J Biol Chem*. 2004; 279:55914–55923. [PubMed: 15485826]
- Shapiro L, Fannon AM, Kwong PD, Thompson A, Lehmann MS, Grubel G, et al. Structural Basis of Cell-Cell Adhesion by Cadherins. *Nature*. 1995; 374:327–337. [PubMed: 7885471]
- Shi QM, Chien YH, Leckband D. Biophysical properties of cadherin bonds do not predict cell sorting. *J Biol Chem*. 2008; 283:28454–28463. [PubMed: 18552401]
- Shi QM, Maruthamuthu V, Li F, Leckband D. Allosteric Cross Talk between Cadherin Extracellular Domains. *Biophys J*. 2010; 99:95–104. [PubMed: 20655837]
- Sivasankar S, Gumbiner B, Leckband D. Direct measurements of multiple adhesive alignments and unbinding trajectories between cadherin extracellular domains. *Biophys J*. 2001; 80:1758–1768. [PubMed: 11259289]

- Sivasankar S, Zhang Y, Nelson WJ, Chu S. Characterizing the initial encounter complex in cadherin adhesion. *Structure*. 2009; 17:1075–1081. [PubMed: 19646884]
- Takeda H, Shimoyama Y, Nagafuchi A, Hirohashi S. E-cadherin functions as a cis-dimer at the cell-cell adhesive interface in vivo. *Nat Struct Biol*. 1999; 6:310–312. [PubMed: 10201395]
- Takeichi M. The Cadherins: cell-cell adhesion molecules controlling animal morphogenesis. *Development*. 1988; 102:639. -&. [PubMed: 3048970]
- Takeichi M. The cadherin superfamily in neuronal connections and interactions. *Nat Rev Neurosci*. 2007; 8:11–20. [PubMed: 17133224]
- Tamura K, Shan WS, Hendrickson WA, Colman DR, Shapiro L. Structure-function analysis of cell adhesion by neural (N-) cadherin. *Neuron*. 1998; 20:1153–1163. [PubMed: 9655503]
- Thomas WE, Trintchina E, Forero M, Vogel V, Sokurenko EV. Bacterial adhesion to target cells enhanced by shear force. *Cell*. 2002; 109:913–923. [PubMed: 12110187]
- Tinkle CL, Lechler T, Pasolli HA, Fuchs E. Conditional targeting of E-cadherin in skin: Insights into hyperproliferative and degenerative responses. *Proc Natl Acad Sci U S A*. 2004; 101:552–557. [PubMed: 14704278]
- Trojanovsky RB, Sokolov E, Trojanovsky SM. Adhesive and lateral E-cadherin dimers are mediated by the same interface. *Mol Cell Biol*. 2003; 23:7965–7972. [PubMed: 14585958]
- Vasioukhin V, Bauer C, Degenstein L, Wise B, Fuchs E. Hyperproliferation and defects in epithelial polarity upon conditional ablation of alpha-catenin in skin. *Cell*. 2001; 104:605–617. [PubMed: 11239416]
- Vendome J, Posy S, Jin X, Bahna F, Ahlsen G, Shapiro L, et al. Molecular design principles underlying Beta-strand swapping in the adhesive dimerization of cadherins. *Nat Struct Mol Biol*. 2011; 18:693–700. [PubMed: 21572446]
- Weber GF, Bjerke MA, DeSimone DW. A Mechanoresponsive Cadherin-Keratin Complex Directs Polarized Protrusive Behavior and Collective Cell Migration. *Dev Cell*. 2012; 22:104–115. [PubMed: 22169071]
- Wu Y, Vendome J, Shapiro L, Ben-Shaul A, Honig B. Transforming binding affinities from three dimensions to two with application to cadherin clustering. *Nature*. 2011; 475:510–513. [PubMed: 21796210]
- Wu YH, Jin XS, Harrison O, Shapiro L, Honig BH, Ben-Shaul A. Cooperativity between trans and cis interactions in cadherin-mediated junction formation. *Proc Natl Acad Sci U S A*. 2010; 107:17592–17597. [PubMed: 20876147]
- Yago T, Wu JH, Wey CD, Klopocki AG, Zhu C, McEver RP. Catch bonds govern adhesion through L-selectin at threshold shear. *J Cell Biol*. 2004; 166:913–923. [PubMed: 15364963]
- Young P, Boussadia O, Halfter H, Grose R, Berger P, Leone DP, et al. E-cadherin controls adherens junctions in the epidermis and the renewal of hair follicles. *EMBO J*. 2003; 22:5723–5733. [PubMed: 14592971]
- Zhang Y, Sivasankar S, Nelson WJ, Chu S. Resolving cadherin interactions and binding cooperativity at the single molecule level. *Proc Natl Acad Sci U S A*. 2009; 106:109–114. [PubMed: 19114658]
- Zhu B, Chappuis-Flament S, Wong E, Jensen IE, Gumbiner BM, Leckband D. Functional analysis of the structural basis of homophilic cadherin adhesion. *Biophys J*. 2003; 84:4033–4042. [PubMed: 12770907]




Figure 1. Adhesive states of classical cadherin and the pathway for cadherin binding
(A) The extracellular region of type-I classical cadherin is composed of five tandem Extracellular (EC) domains. Linkers between successive EC domains are each bound to three Ca^{2+} ions which give the ectodomain its characteristic curvature. Ectodomains from opposing cells (shown in green and magenta) adhere across the inter-membrane gap via ‘*trans*’ interactions. The primary *trans* interface involves the interaction of opposing EC1 domains and is termed the strand-swapped dimer. In this conformation, N-terminal β -strands between opposing EC1 domains are swapped and the side chain of a conserved Tryptophan at position 2 (W2) is inserted into a pocket on their adhesive partner. **(B)** Prior to strand-swapping, cadherin ectodomains form a non-swapped, intermediate conformation, called an X-dimer. This conformation is formed by extensive surface interactions between the base of the EC1 domain, EC1–EC2 inter-domain linker region and the apex of domain EC2. **(C)** Cadherin monomers adopt a “closed” conformation where W2 is docked into each monomer’s binding pocket. Monomers from opposing cells interact to form X-dimers and then proceed to swap W2 residues to form a strand-swap dimer. The K_d of the EC1–2

domains of W2A E-cadherin X-dimers is 916 μM (Harrison *et al.*, 2010) while the K_d of the EC1–2 domains of WT E-cadherin strand-swap dimers is 97 μM (Katsamba *et al.*, 2009).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2. Mechanical force tunes the kinetics of cadherin adhesion

Adapted from (Rakshit et al., 2012). (A) X-dimers form catch bonds which become longer lived and lock in the presence of tensile force. When W2A cadherin X-dimers are pulled, their bond lifetimes increase with force. After reaching a maximum at a critical force of ~ 30 pN, the lifetimes decrease. (B) Strand swap dimers form slip bonds which become shorter lived when pulled. Slip bonds are formed by K14E mutants that interact for short and long periods of time and also by WT cadherins that interact for long periods of time. However, when WT cadherins interact for a short period of time, they form ideal bonds that are insensitive to force. (C) Hypothetical mechanism by which keratinocytes resist tensile forces during skin renewal and wound healing. As skin cells reposition themselves, E-cadherins bind rapidly to form X-dimers that allow cells to grip strongly under load. In immobile keratinocytes, E-cadherins form more robust strand-swap dimers that have a high affinity in the absence of force.